
Data is for the week ending 12/4/2010 (MMWR week 48) Created 12/8/2010


Influenza-like Illness Activity



Influenza-like Illness (ILI) Activity Levels by Health District - Utah

| IZA-IIKC IIIIIC33 (ILI) ACTIVI | ity Ecvels by ricaltii District | | |
|--------------------------------|---------------------------------|--|--|
| Health District | ILI Activity | | |
| Bear River | Minimal | | |
| Central | No Data | | |
| Davis | Minimal | | |
| Salt Lake | Minimal | | |
| Southwest | No Data | | |
| Summit | Minimal | | |
| Tooele | Minimal | | |
| TriCounty | No Data | | |
| Utah | Minimal | | |
| Wasatch | Minimal | | |
| Weber-Morgan | No Data | | |
| State | Minimal | | |

Severity Measures



^{*}Event date is calculated based on a hierarchy of dates: 1. onset date 2. specimen collection date 3. date reported to public health.

Summary Data for Influenza-associated Hospitalizations, 2010-2011 Season to Date

| Variable | | Number of | % of IAH | % in Utah | p value* |
|----------|----------------------------------|-----------|----------|-----------|----------|
| | | cases | cases | рор | p value |
| | Male | 5 | 22.7% | 50.3% | 0.0097 |
| Sex | Female | 17 | 77.3% | 49.7% | 0.0097 |
| | Unknown | 0 | 0.0% | NA | |
| | White, Not Hispanic | 9 | 40.9% | 82.0% | < 0.0001 |
| | Hispanic | 10 | 45.5% | 11.6% | < 0.0001 |
| | Native Hawaiian/Pacific Islander | 3 | 13.6% | 0.7% | < 0.0001 |
| Race | Black/African American | 0 | 0.0% | 0.9% | 0.6490 |
| America | American Indian | 0 | 0.0% | 1.1% | 0.6157 |
| | Asian | 0 | 0.0% | 1.9% | 0.5174 |
| | Unknown | 0 | 0.0% | NA | |

^{*}If a p value is \le 0.05, there is a significant difference between the percentage seen in IAH and the general Utah population.

Data is for the week ending 12/4/2010 (MMWR week 48, Created 12/8/2010

Severity Measures, cont.

Cases of Influenza-associated Hospitalizations by Age Group, Utah, 2010-2011

| Age Group | Total Cases Season to Date | Percent of Cases | Cases/ 100,000 Population | New Cases in Past Week | |
|-------------|----------------------------------|------------------|---------------------------------|------------------------------|--|
| 0-4 years | 6 | 27.3% | 2.2 | 0 | |
| 5-24 years | 4 | 18.2% | 0.4 | 0 | |
| 25-49 years | 3 | 13.6% | 0.3 | 0 | |
| 50-64 years | 5 | 22.7% | 1.2 | 1 | |
| 65+ years | 4 | 18.2% | 1.6 | 0 | |
| Total | 22 | 100% | 0.8 | 1 | |

Influenza-associated Hospitalizations by Health District, 2010-2011 Season to Date

| Health District | Total Hospitalizations |
|-----------------|------------------------|
| Bear River | 0 |
| Central | 0 |
| Davis | 0 |
| Salt Lake | 18 |
| Southeastern | 0 |
| Southwest | 0 |
| Summit | 1 |
| Tooele | 1 |
| TriCounty | 0 |
| Utah | 0 |
| Wasatch | 0 |
| Weber-Morgan | 2 |

Summary Data for Influenza-associated Hospitalizations, 2010-2011 Season to Date

| • | Yes | | No | | Unknown | |
|------------------------------|--------------------|------------|--------------------|------------|-----------------|------------|
| Variable | Number of Cases | % of cases | Number of Cases | % of cases | Number of Cases | % of cases |
| ICU | 2 | 9.1% | 10 | 45.5% | 10 | 45.5% |
| Ventilator | 1 | 4.5% | 11 | 50.0% | 10 | 45.5% |
| Died | 0 | 0.0% | 12 | 54.5% | 10 | 45.5% |
| Neurological Symptoms | 1 | 4.5% | 10 | 45.5% | 11 | 50.0% |
| Healthcare Worker | 0 | 0.0% | 12 | 54.5% | 10 | 45.5% |
| Pregnant | 0 | 0.0% | 17 | 77.3% | 5 | 22.7% |
| Heart Disorder | 0 | 0.0% | 11 | 50.0% | 11 | 50.0% |
| Blood Disorder | 0 | 0.0% | 11 | 50.0% | 11 | 50.0% |
| Kidney Disorder | 0 | 0.0% | 11 | 50.0% | 11 | 50.0% |
| Metabolic Disorder | 1 | 4.5% | 10 | 45.5% | 11 | 50.0% |
| Chronic Respiratory Disorder | 4 | 18.2% | 7 | 31.8% | 11 | 50.0% |
| Immunosuppressed | 2 | 9.1% | 9 | 40.9% | 11 | 50.0% |
| Neurological Disorder | 1 | 4.5% | 10 | 45.5% | 11 | 50.0% |
| Seizure Disorder | 0 | 0.0% | 11 | 50.0% | 11 | 50.0% |
| Bacterial Co-infection | 0 | 0.0% | 11 | 50.0% | 11 | 50.0% |
| Obese** | 2 | 15.4% | 4 | 30.8% | 7 | 53.8% |
| Morbidly Obese** | 2 | 15.4% | 4 | 30.8% | 7 | 53.8% |
| Risk Factor*** | 20 | 90.9% | 2 | 9.1% | 0 | 0.0% |
| Vaccinated | 2 | 9.1% | 8 | 36.4% | 12 | 54.5% |

^{**}Obesity and morbid obesity is not considered for individuals under 18 years or pregnant women. Thus total counts will not equal the total number of influenza-associated hospitalizations

Case Counts of Influenza-associated Hospitalizations, Utah, 2010-2011

| Case Status | Current Week | Current Week % of Cases | Season Total | Season % of Cases |
|-------------|-----------------|----------------------------|-----------------|-------------------|
| Confirmed* | 1 | 100.0% | 21 | 95.5% |
| Probable** | 0 | 0.0% | 1 | 4.5% |
| Total | 1 | 100.0% | 22 | 100.0% |


^{*}Confirmed= hospitalized with serology, DFA, PCR, or culture positive influenza test

^{***}Risk factors for influenza include: persons < 5 years, persons ≥ 65 years, pregnant women, and persons with a chronic medical condition.


^{**}Probable= hospitalized with a positive rapid test

Data is for the week ending 12/4/2010 (MMWR week 48) Created 12/8/2010

Pneumonia and Influenza Deaths



School Absenteeism



Weekly Student Absenteeism - Utah, 2010-2011

| Health District | All absences per 100 students/day | | | |
|-----------------|-----------------------------------|--|--|--|
| Bear River | 3.2 | | | |
| Central | 3.4 | | | |
| Davis | 2.4 | | | |
| Salt Lake | 4.7 | | | |
| Southeast | 5.2 | | | |
| Southwest | | | | |
| Summit | | | | |
| Tooele | 6.2 | | | |
| TriCounty | 8.3 | | | |
| Utah | 5.2 | | | |
| Wasatch | 6.0 | | | |
| Weber-Morgan | 6.4 | | | |
| State | 5.2 | | | |

Data is for the week ending 12/4/2010 (MMWR week 48) Created 12/8/2010

Virologic Data

Unified State Laboratory: Public Health Virologic Data

| | Season to-date | | Current MMWR weel | |
|-----------------------------|----------------|-------------|-------------------|---------|
| | Number | Percent | Number | Percent |
| Specimens tested | 184 | - | 33 | |
| Positive specimens | 74 | 40.2% | 19 | 57.6% |
| Positive Spe | cimens by Ty | /pe/Subtype | | |
| Influenza A | 73 | 98.6% | 19 | 100.0% |
| A (2009 H1N1) | 3 | 4.1% | 0 | 0.0% |
| A (H1) | 0 | 0.0% | 0 | 0.0% |
| A (H3) | 70 | 95.9% | 19 | 100.0% |
| A (subtyping not performed) | 0 | 0.0% | 0 | 0.0% |
| A (unable to subtype) | 0 | 0.0% | 0 | 0.0% |
| Influenza B | 2 | 2.7% | 0 | 0.0% |

